



190.28 - Tumor Antigen by Immunoassay CA 125

Description

Immunoassay determinations of the serum levels of certain proteins or carbohydrates serve as tumor markers. When elevated, serum concentration of these markers may reflect tumor size and grade. This policy specifically addresses tumor antigen CA 125.

HCPCS Codes (Alphanumeric, CPT® AMA)

Code	Description
86304	Immunoassay for tumor antigen, quantitative, CA 125

ICD-10-CM Codes Covered by Medicare Program

The ICD-10-CM codes in the table below can be viewed on CMS' website as part of Downloads: Lab Code List, at
<http://www.cms.gov/Medicare/Coverage/CoverageGenInfo/LabNCDsICD10.html>

Code	Description
C45.1	Mesothelioma of peritoneum
C48.1	Malignant neoplasm of specified parts of peritoneum
C48.2	Malignant neoplasm of peritoneum, unspecified
C48.8	Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum
C51.8	Malignant neoplasm of overlapping sites of vulva
C53.0	Malignant neoplasm of endocervix
C54.1	Malignant neoplasm of endometrium
C54.2	Malignant neoplasm of myometrium
C54.3	Malignant neoplasm of fundus uteri
C54.9	Malignant neoplasm of corpus uteri, unspecified
C56.1	Malignant neoplasm of right ovary
C56.2	Malignant neoplasm of left ovary
C56.3	Malignant neoplasm of bilateral ovaries
C56.9	Malignant neoplasm of unspecified ovary
C57.00	Malignant neoplasm of unspecified fallopian tube
C57.01	Malignant neoplasm of right fallopian tube
C57.02	Malignant neoplasm of left fallopian tube
C57.4	Malignant neoplasm of uterine adnexa, unspecified
C57.7	Malignant neoplasm of other specified female genital organs
C57.8	Malignant neoplasm of overlapping sites of female genital organs



**Medicare National Coverage Determinations (NCD)
Coding Policy Manual and Change Report (ICD-10-CM)**

Code	Description
C79.60	Secondary malignant neoplasm of unspecified ovary
C79.61	Secondary malignant neoplasm of right ovary
C79.62	Secondary malignant neoplasm of left ovary
C79.63	Secondary malignant neoplasm of bilateral ovaries
C79.82	Secondary malignant neoplasm of genital organs
D39.0	Neoplasm of uncertain behavior of uterus
D39.10	Neoplasm of uncertain behavior of unspecified ovary
D39.11	Neoplasm of uncertain behavior of right ovary
D39.12	Neoplasm of uncertain behavior of left ovary
D39.2	Neoplasm of uncertain behavior of placenta
D39.8	Neoplasm of uncertain behavior of other specified female genital organs
D39.9	Neoplasm of uncertain behavior of female genital organ, unspecified
G89.3	Neoplasm related pain (acute) (chronic)
M33.03	Juvenile dermatomyositis without myopathy
M33.13	Other dermatomyositis without myopathy
M33.93	Dermatopolymyositis, unspecified without myopathy
R19.01	Right upper quadrant abdominal swelling, mass and lump
R19.02	Left upper quadrant abdominal swelling, mass and lump
R19.03	Right lower quadrant abdominal swelling, mass and lump
R19.04	Left lower quadrant abdominal swelling, mass and lump
R19.09	Other intra-abdominal and pelvic swelling, mass and lump
R97.1	Elevated cancer antigen 125 [CA 125]
R97.8	Other abnormal tumor markers
Z85.41	Personal history of malignant neoplasm of cervix uteri
Z85.42	Personal history of malignant neoplasm of other parts of uterus
Z85.43	Personal history of malignant neoplasm of ovary
Z85.44	Personal history of malignant neoplasm of other female genital organs

Indications

CA 125 is a high molecular weight serum tumor marker elevated in 80% of patients who present with epithelial ovarian carcinoma. It is also elevated in carcinomas of the fallopian tube, endometrium, and endocervix. An elevated level may also be associated with the presence of a malignant mesothelioma or primary peritoneal carcinoma.



A CA 125 level may be obtained as part of the initial pre-operative work-up for women presenting with a suspicious pelvic mass to be used as a baseline for purposes of post-operative monitoring. Initial declines in CA 125 after initial surgery and/or chemotherapy for ovarian carcinoma are also measured by obtaining three serum levels during the first month post treatment to determine the patient's CA 125 half-life, which has significant prognostic implications.

The CA 125 levels are again obtained at the completion of chemotherapy as an index of residual disease. Surveillance CA 125 measurements are generally obtained every 3 months for 2 years, every 6 months for the next 3 years, and yearly thereafter. CA 125 levels are also an important indicator of a patient's response to therapy in the presence of advanced or recurrent disease. In this setting, CA 125 levels may be obtained prior to each treatment cycle.

Limitations

These services are not covered for the evaluation of patients with signs or symptoms suggestive of malignancy. The service may be ordered at times necessary to assess either the presence of recurrent disease or the patient's response to treatment with subsequent treatment cycles.

The CA 125 is specifically not covered for aiding in the differential diagnosis of patients with a pelvic mass as the sensitivity and specificity of the test is not sufficient. In general, a single "tumor marker" will suffice in following a patient with one of these malignancies.

ICD-10-CM Codes That Do Not Support Medical Necessity

Any ICD-10-CM code not listed in either of the ICD-10-CM covered or non-covered sections.

Documentation Requirements

Indicated if service request for CA125 is requested more frequently than stipulated.

Sources of Information

Clinical Pancreatic Guideline for the Use of Tumor Markers in Breast and Colorectal Cancer, American Society of Clinical Oncology. J Clin Oncol 14:2843-2877, 1996.

Chan DW, Beveridge RA, Muss H, et al. Use of Triquant BR Radioimmunoassay for Early Detection of Breast Cancer Recurrence in Patients with Stage II and Stage III Disease. J Clin Oncol 1977, 15(6):2322-2328.



190.29 - Tumor Antigen by Immunoassay CA 15-3/CA 27.29

Description

Immunoassay determinations of the serum levels of certain proteins or carbohydrates serve as tumor markers. When elevated, serum concentration of markers may reflect tumor size & grade. This policy specifically addresses the following tumor antigens: CA 15-3 and CA 27.29

HCPCS Codes (Alphanumeric, CPT® AMA)

Code	Description
86300	Immunoassay for tumor antigen, quantitative; CA 15-3 (27.29)

ICD-10-CM Codes Covered by Medicare Program

The ICD-10-CM codes in the table below can be viewed on CMS' website as part of Downloads: Lab Code List, at
<http://www.cms.gov/Medicare/Coverage/CoverageGenInfo/LabNCDsICD10.html>

Code	Description
C44.1321	Sebaceous cell carcinoma of skin of right upper eyelid, including canthus
C44.1322	Sebaceous cell carcinoma of skin of right lower eyelid, including canthus
C44.1391	Sebaceous cell carcinoma of skin of left upper eyelid, including canthus
C44.1392	Sebaceous cell carcinoma of skin of left lower eyelid, including canthus
C50.011	Malignant neoplasm of nipple and areola, right female breast
C50.012	Malignant neoplasm of nipple and areola, left female breast
C50.019	Malignant neoplasm of nipple and areola, unspecified female breast
C50.021	Malignant neoplasm of nipple and areola, right male breast
C50.022	Malignant neoplasm of nipple and areola, left male breast
C50.029	Malignant neoplasm of nipple and areola, unspecified male breast
C50.111	Malignant neoplasm of central portion of right female breast
C50.112	Malignant neoplasm of central portion of left female breast
C50.119	Malignant neoplasm of central portion of unspecified female breast
C50.121	Malignant neoplasm of central portion of right male breast
C50.122	Malignant neoplasm of central portion of left male breast
C50.129	Malignant neoplasm of central portion of unspecified male breast
C50.211	Malignant neoplasm of upper-inner quadrant of right female breast
C50.212	Malignant neoplasm of upper-inner quadrant of left female breast
C50.219	Malignant neoplasm of upper-inner quadrant of unspecified female breast
C50.221	Malignant neoplasm of upper-inner quadrant of right male breast



**Medicare National Coverage Determinations (NCD)
Coding Policy Manual and Change Report (ICD-10-CM)**

Code	Description
C50.222	Malignant neoplasm of upper-inner quadrant of left male breast
C50.229	Malignant neoplasm of upper-inner quadrant of unspecified male breast
C50.311	Malignant neoplasm of lower-inner quadrant of right female breast
C50.312	Malignant neoplasm of lower-inner quadrant of left female breast
C50.319	Malignant neoplasm of lower-inner quadrant of unspecified female breast
C50.321	Malignant neoplasm of lower-inner quadrant of right male breast
C50.322	Malignant neoplasm of lower-inner quadrant of left male breast
C50.329	Malignant neoplasm of lower-inner quadrant of unspecified male breast
C50.411	Malignant neoplasm of upper-outer quadrant of right female breast
C50.412	Malignant neoplasm of upper-outer quadrant of left female breast
C50.419	Malignant neoplasm of upper-outer quadrant of unspecified female breast
C50.421	Malignant neoplasm of upper-outer quadrant of right male breast
C50.422	Malignant neoplasm of upper-outer quadrant of left male breast
C50.429	Malignant neoplasm of upper-outer quadrant of unspecified male breast
C50.511	Malignant neoplasm of lower-outer quadrant of right female breast
C50.512	Malignant neoplasm of lower-outer quadrant of left female breast
C50.519	Malignant neoplasm of lower-outer quadrant of unspecified female breast
C50.521	Malignant neoplasm of lower-outer quadrant of right male breast
C50.522	Malignant neoplasm of lower-outer quadrant of left male breast
C50.529	Malignant neoplasm of lower-outer quadrant of unspecified male breast
C50.611	Malignant neoplasm of axillary tail of right female breast
C50.612	Malignant neoplasm of axillary tail of left female breast
C50.619	Malignant neoplasm of axillary tail of unspecified female breast
C50.621	Malignant neoplasm of axillary tail of right male breast
C50.622	Malignant neoplasm of axillary tail of left male breast
C50.629	Malignant neoplasm of axillary tail of unspecified male breast
C50.811	Malignant neoplasm of overlapping sites of right female breast
C50.812	Malignant neoplasm of overlapping sites of left female breast
C50.819	Malignant neoplasm of overlapping sites of unspecified female breast
C50.821	Malignant neoplasm of overlapping sites of right male breast
C50.822	Malignant neoplasm of overlapping sites of left male breast
C50.829	Malignant neoplasm of overlapping sites of unspecified male breast
C50.911	Malignant neoplasm of unspecified site of right female breast
C50.912	Malignant neoplasm of unspecified site of left female breast

NCD 190.29

***January 2024 Changes
ICD-10-CM Version – Red**



Code	Description
C50.919	Malignant neoplasm of unspecified site of unspecified female breast
C50.921	Malignant neoplasm of unspecified site of right male breast
C50.922	Malignant neoplasm of unspecified site of left male breast
C50.929	Malignant neoplasm of unspecified site of unspecified male breast
C79.2	Secondary malignant neoplasm of skin
C79.81	Secondary malignant neoplasm of breast
C84.7A	Anaplastic large cell lymphoma, ALK-negative, breast
G89.3	Neoplasm related pain (acute) (chronic)
R97.8	Other abnormal tumor markers
Z85.3	Personal history of malignant neoplasm of breast
Z86.002	Personal history of in-situ neoplasm of other and unspecified genital organs
Z86.003	Personal history of in-situ neoplasm of oral cavity, esophagus and stomach
Z86.004	Personal history of in-situ neoplasm of other and unspecified digestive organs
Z86.005	Personal history of in-situ neoplasm of middle ear and respiratory system
Z86.006	Personal history of melanoma in-situ
Z86.007	Personal history of in-situ neoplasm of skin

Indications

Multiple tumor markers are available for monitoring the response of certain malignancies to therapy and assessing whether a residual tumor exists post-surgical therapy. CA 15-3 is often medically necessary to aid in the management of patients with breast cancer. Serial testing must be used in conjunction with other clinical methods for monitoring breast cancer. For monitoring, if medically necessary, use consistently either CA 15-3 or CA 27.29, not both. CA 27.29 is equivalent to CA 15-3 in its usage in management of patients with breast cancer.

Limitations

These services are not covered for the evaluation of patients with signs or symptoms suggestive of malignancy. The service may be ordered at times necessary to assess either the presence of recurrent disease or the patient’s response to treatment with subsequent treatment cycles.

ICD-10-CM Codes That Do Not Support Medical Necessity

Any ICD-10-CM code not listed in either of the ICD-10-CM covered or non-covered sections.

Sources of Information

Clinical Pancreatic Guideline for the Use of Tumor Markers in Breast and Colorectal Cancer, American Society of Clinical Oncology. J Clin Oncol 14:2843-2877, 1996.



**Medicare National Coverage Determinations (NCD)
Coding Policy Manual and Change Report (ICD-10-CM)**

Chan DW, Beveridge RA, Muss H, et al. Use of Triquant BR Radioimmunoassay for Early Detection of Breast Cancer Recurrence in Patients with Stage II & Stage III Disease. *J Clin Oncol* 1977, 15(6):2322-2328.

Bone GG, von Mensdorff-Pouilly S, Kenemans P, van Kamp GJ, et al. Clinical and Technical Evaluation of ACS BR Serum Assay of MUC-1 Gene Derived Glycoprotein in Breast Cancer, and Compared with CA15-3 Assays. *Clin Chem* 1997, 43(4):585-593.