

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

Healthy Michigan Plan

The Patient Protection and Affordable Care Act have designated specific resources that identify the preventive services required for coverage by the act. The following lists of preventive services, CPT, and HCPCS codes are for reference purposes only to assist Medicaid managed care plans in determining coverage for specific services as required for Healthy Michigan Plan beneficiaries. Other coverage determination guidelines may apply. This information is not intended to replace clinical judgment of primary care providers in administering these services.

Additional information about preventive care guidelines is available at the following:

- United States Preventive Services Task Force grade A and B services
<http://www.uspreventiveservicestaskforce.org/Page/Name/recommendations>
- Advisory Committee on Immunization Practices recommended vaccines
<http://www.cdc.gov/vaccines/hcp/acip-recs/index.html>
- Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine recommended preventive services for women
<http://www.nationalacademies.org/hmd/Reports/2011/Clinical-Preventive-Services-for-Women-Closing-the-Gaps.aspx>
- Early and Periodic Screening, Diagnosis and Treatment services as defined in the current periodicity schedule by the American Academy of Pediatrics (for 19 and 20 year olds only)
http://brightfutures.aap.org/clinical_practice.html

USPSTF A and B RECOMMENDATIONS

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Abdominal aortic aneurysm screening: men	The USPSTF recommends one-time screening for abdominal aortic aneurysm by ultrasonography in men ages 65 to 75 years who have ever smoked.	N/A	Healthy Michigan Plan does not cover this age group.
Alcohol misuse: screening and counseling	The USPSTF recommends that clinicians screen adults age 18 years or older for alcohol misuse and provide persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce alcohol misuse.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Aspirin preventive medication: adults aged 50 to 59 years with a ≥10% 10-year cardiovascular risk	The USPSTF recommends initiating low-dose aspirin use for the primary prevention of cardiovascular disease and colorectal cancer in adults aged 50 to 59 years who have a 10% or greater 10-year cardiovascular risk, are not at increased risk for bleeding, have a life expectancy of at least 10 years, and are willing to take low-dose aspirin daily for at least 10 years.	N/A	Pharmacy Benefit
Bacteriuria screening: pregnant women	The USPSTF recommends screening for asymptomatic bacteriuria with urine culture in pregnant women at 12 to 16 weeks' gestation or at the first prenatal visit, if later.	87086, 87088	

Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines
Healthy Michigan Plan

USPSTF A and B RECOMMENDATIONS

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Blood pressure screening in adults	The USPSTF recommends screening for high blood pressure in adults aged 18 years or older. The USPSTF recommends obtaining measurements outside of the clinical setting for diagnostic confirmation before starting treatment.	See Notes	Covered during preventive medicine services or a focused E/M visit.
BRCA risk assessment and genetic counseling/testing	The USPSTF recommends that primary care providers screen women who have family members with breast, ovarian, tubal, or peritoneal cancer with one of several screening tools designed to identify a family history that may be associated with an increased risk for potentially harmful mutations in breast cancer susceptibility genes (BRCA1 or BRCA2). Women with positive screening results should receive genetic counseling and, if indicated after counseling, BRCA testing.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Breast cancer preventive medications	The USPSTF recommends that clinicians engage in shared, informed decision making with women who are at increased risk for breast cancer about medications to reduce their risk. For women who are at increased risk for breast cancer and at low risk for adverse medication effects, clinicians should offer to prescribe risk-reducing medications, such as tamoxifen or raloxifene.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Breast cancer screening	The USPSTF recommends screening mammography for women, with or without clinical breast examination, every 1 to 2 years for women age 40 years and older.	77067, 77063, G0202	
Breastfeeding interventions	The USPSTF recommends providing interventions during pregnancy and after birth to support breastfeeding.	99402-MIHP S9443 Also See Notes	Lactation support and counseling is provided as part of childbirth education. It is covered during preventive medicine services or a focused E/M visit. IBCLC services may be provided as a benefit within the MIHP program or separately reimbursed in the outpatient setting when rendered by a qualified provider

Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines
Healthy Michigan Plan

USPSTF A and B RECOMMENDATIONS

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Cervical cancer screening	The USPSTF recommends screening for cervical cancer in women ages 21 to 65 years with cytology (Pap smear) every 3 years or, for women ages 30 to 65 years who want to lengthen the screening interval, screening with a combination of cytology and human papillomavirus (HPV) testing every 5 years.	87623-87625, 88141-88148, 88155, 88164-88167, 88174-88175, G0101, G0476, Q0091	
Chlamydia screening: women	The USPSTF recommends screening for chlamydia in sexually active women age 24 years or younger and in older women who are at increased risk for infection.	87110, 87270, 87320, 87490-87492, 87810	
Cholesterol abnormalities screening: men 35 and older	The USPSTF strongly recommends screening men age 35 years and older for lipid disorders.	80061, 82465, 83718-83721, 84478	
Cholesterol abnormalities screening: men younger than 35	The USPSTF recommends screening men ages 20 to 35 years for lipid disorders if they are at increased risk for coronary heart disease.	80061, 82465, 83718-83721, 84478	
Cholesterol abnormalities screening: women 45 and older	The USPSTF strongly recommends screening women age 45 years and older for lipid disorders if they are at increased risk for coronary heart disease.	80061, 82465, 83718-83721, 84478	
Cholesterol abnormalities screening: women younger than 45	The USPSTF recommends screening women ages 20 to 45 years for lipid disorders if they are at increased risk for coronary heart disease.	80061, 82465, 83718-83721, 84478	
Colorectal cancer screening	The USPSTF recommends screening for colorectal cancer starting at age 50 years and continuing until age 75 years.	45330-45331, 45333, 45338, 45346, 45378, 45380, 45384-45385, 45388, 81528, 82270, 82274, G0104-G0105, G0121, G0328	
Dental caries prevention: infants and children up to age 5 years	The USPSTF recommends the application of fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption in primary care practices. The USPSTF recommends primary care clinicians prescribe oral fluoride supplementation starting at age 6 months for children whose water supply is fluoride deficient.	N/A	Healthy Michigan Plan does not cover this age group.

Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines
Healthy Michigan Plan

USPSTF A and B RECOMMENDATIONS

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Depression screening: adolescents	The USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12 to 18 years. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.	N/A	Healthy Michigan Plan does not cover this age group.
Depression screening: adults	The USPSTF recommends screening for depression in the general adult population, including pregnant and postpartum women. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Diabetes screening	The USPSTF recommends screening for abnormal blood glucose as part of cardiovascular risk assessment in adults aged 40 to 70 years who are overweight or obese. Clinicians should offer or refer patients with abnormal blood glucose to intensive behavioral counseling interventions to promote a healthful diet and physical activity.	82947-82952, 83036	
Falls prevention in older adults: exercise or physical therapy	The USPSTF recommends exercise or physical therapy to prevent falls in community-dwelling adults age 65 years and older who are at increased risk for falls.	N/A	Healthy Michigan Plan does not cover this age group.
Falls prevention in older adults: vitamin D	The USPSTF recommends vitamin D supplementation to prevent falls in community-dwelling adults age 65 years and older who are at increased risk for falls.	N/A	Healthy Michigan Plan does not cover this age group.
Folic acid supplementation	The USPSTF recommends that all women planning or capable of pregnancy take a daily supplement containing 0.4 to 0.8 mg (400 to 800 µg) of folic acid.	N/A	Pharmacy Benefit
Gestational diabetes mellitus screening	The USPSTF recommends screening for gestational diabetes mellitus in asymptomatic pregnant women after 24 weeks of gestation.	82950-82952	
Gonorrhea prophylactic medication: newborns	The USPSTF recommends prophylactic ocular topical medication for all newborns for the prevention of gonococcal ophthalmia neonatorum.	N/A	Healthy Michigan Plan does not cover this age group.
Gonorrhea screening: women	The USPSTF recommends screening for gonorrhea in sexually active women age 24 years or younger and in older women who are at increased risk for infection.	87590-87592, 87850	

Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines
Healthy Michigan Plan

USPSTF A and B RECOMMENDATIONS

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Healthy diet and physical activity counseling to prevent cardiovascular disease: adults with cardiovascular risk factors	The USPSTF recommends offering or referring adults who are overweight or obese and have additional cardiovascular disease (CVD) risk factors to intensive behavioral counseling interventions to promote a healthful diet and physical activity for CVD prevention.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Hearing loss screening: newborns	The USPSTF recommends screening for hearing loss in all newborn infants.	N/A	Healthy Michigan Plan does not cover this age group.
Hemoglobinopathies screening: newborns	The USPSTF recommends screening for sickle cell disease in newborns.	N/A	Healthy Michigan Plan does not cover this age group.
Hepatitis B screening: nonpregnant adolescents and adults	The USPSTF recommends screening for hepatitis B virus infection in persons at high risk for infection.	86704-86706, 87340, G0499	
Hepatitis B screening: pregnant women	The USPSTF strongly recommends screening for hepatitis B virus infection in pregnant women at their first prenatal visit.	86704-86706, 87340, G0499	
Hepatitis C virus infection screening: adults	The USPSTF recommends screening for hepatitis C virus (HCV) infection in persons at high risk for infection. The USPSTF also recommends offering one-time screening for HCV infection to adults born between 1945 and 1965.	86803, G0472	
High blood pressure in adults: screening	The USPSTF recommends screening for high blood pressure in adults aged 18 years or older. The USPSTF recommends obtaining measurements outside of the clinical setting for diagnostic confirmation before starting treatment.	See Notes	Covered during preventive medicine services or a focused E/M visit.
HIV screening: nonpregnant adolescents and adults	The USPSTF recommends that clinicians screen for HIV infection in adolescents and adults ages 15 to 65 years. Younger adolescents and older adults who are at increased risk should also be screened.	86689, 86701-86703, 87389-87391, 87534-87539, 87806, G0432-G0433, G0435	

Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines
Healthy Michigan Plan

USPSTF A and B RECOMMENDATIONS

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
HIV screening: pregnant women	The USPSTF recommends that clinicians screen all pregnant women for HIV, including those who present in labor who are untested and whose HIV status is unknown.	80081, 86689, 86701-86703, 87389-87391, 87534-87539, 87806, G0432-G0433, G0435	
Hypothyroidism screening: newborns	The USPSTF recommends screening for congenital hypothyroidism in newborns.	N/A	Healthy Michigan Plan does not cover this age group.
Intimate partner violence screening: women of childbearing age	The USPSTF recommends that clinicians screen women of childbearing age for intimate partner violence, such as domestic violence, and provide or refer women who screen positive to intervention services. This recommendation applies to women who do not have signs or symptoms of abuse.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Lung cancer screening	The USPSTF recommends annual screening for lung cancer with low-dose computed tomography in adults' ages 55 to 80 years who have a 30 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.	71250, G0297	
Obesity screening and counseling: adults	The USPSTF recommends screening all adults for obesity. Clinicians should offer or refer patients with a body mass index of 30 kg/m2 or higher to intensive, multicomponent behavioral interventions.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Obesity screening and counseling: children	The USPSTF recommends that clinicians screen children age 6 years and older for obesity and offer them or refer them to comprehensive, intensive behavioral interventions to promote improvement in weight status.	N/A	Healthy Michigan Plan does not cover this age group.
Osteoporosis screening: women	The USPSTF recommends screening for osteoporosis in women age 65 years and older and in younger women whose fracture risk is equal to or greater than that of a 65-year-old white woman who has no additional risk factors.	76977, 77078-77081	
Phenylketonuria screening: newborns	The USPSTF recommends screening for phenylketonuria in newborns.	N/A	Healthy Michigan Plan does not cover this age group.

Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines
Healthy Michigan Plan

USPSTF A and B RECOMMENDATIONS

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Preeclampsia prevention: aspirin	The USPSTF recommends the use of low-dose aspirin (81 mg/d) as preventive medication after 12 weeks of gestation in women who are at high risk for preeclampsia.	N/A	Pharmacy Benefit
Rh incompatibility screening: 24–28 weeks' gestation	The USPSTF recommends repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24 to 28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.	80055, 80081, 86901	
Rh incompatibility screening: first pregnancy visit	The USPSTF strongly recommends Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.	80055, 80081, 86901	
Sexually transmitted infections counseling	The USPSTF recommends intensive behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Skin cancer behavioral counseling	The USPSTF recommends counseling children, adolescents, and young adults' ages 10 to 24 years who have fair skin about minimizing their exposure to ultraviolet radiation to reduce risk for skin cancer.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Statin preventive medication: adults ages 40–75 years with no history of CVD, 1 or more CVD risk factors, and a calculated 10-year CVD event risk of 10% or greater	The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (i.e., symptomatic coronary artery disease or ischemic stroke) use a low- to moderate-dose statin for the prevention of CVD events and mortality when all of the following criteria are met: 1) they are ages 40 to 75 years; 2) they have 1 or more CVD risk factors (i.e., dyslipidemia, diabetes, hypertension, or smoking); and 3) they have a calculated 10-year risk of a cardiovascular event of 10% or greater. Identification of dyslipidemia and calculation of 10-year CVD event risk requires universal lipids screening in adults ages 40 to 75 years.	N/A	Pharmacy Benefit
Syphilis screening: nonpregnant persons	The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection.	86592-86593	
Syphilis screening: pregnant women	The USPSTF recommends that clinicians screen all pregnant women for syphilis infection.	80055, 80081, 86592-86593	

Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines
Healthy Michigan Plan

USPSTF A and B RECOMMENDATIONS

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Tobacco use counseling and interventions: nonpregnant adults	The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and U.S. Food and Drug Administration (FDA)–approved pharmacotherapy for cessation to adults who use tobacco.	99406, 99407	
Tobacco use counseling: pregnant women	The USPSTF recommends that clinicians ask all pregnant women about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant women who use tobacco.	99406, 99407	
Tobacco use interventions: children and adolescents	The USPSTF recommends that clinicians provide interventions, including education or brief counseling, to prevent initiation of tobacco use in school-aged children and adolescents.	N/A	Healthy Michigan Plan does not cover this age group.
Tuberculosis screening: adults	The USPSTF recommends screening for latent tuberculosis infection in populations at increased risk.	86480, 86481, 86580, 87116	
Visual acuity screening in children	The USPSTF recommends vision screening for all children at least once between the ages of 3 and 5 years, to detect the presence of amblyopia or its risk factors.	N/A	Healthy Michigan Plan does not cover this age group.

Advisory Committee on Immunization Practices (ACIP) recommended vaccines

Follow current Medicaid policy for coverage of ACIP recommended vaccines.

Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine
Recommended Preventive Health Care Services for Women

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Screening for gestational diabetes (Recommendation 5.1)	Screening for gestational diabetes in pregnant women between 24 and 28 weeks of gestation and at the first prenatal visit for pregnant women identified to be at high risk for diabetes.	82950-82952	

Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines
Healthy Michigan Plan

Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine

Recommended Preventive Health Care Services for Women

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Human papillomavirus testing (Recommendation 5.2)	The addition of high-risk human papillomavirus DNA testing in addition to cytology testing in women with normal cytology results. Screening should begin at 30 years of age and should occur no more frequently than every 3 years.	87623-87625, G0476	
Counseling for sexually transmitted infections (Recommendation 5.3)	Annual counseling on sexually transmitted infections for sexually active women.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Counseling and screening for human immune-deficiency virus (Recommendation 5.4)	Counseling and screening for human immunodeficiency virus infection on an annual basis for sexually active women.	86689, 86701-86703, 87389-87391, 87534-87539, 87806, G0432-G0433, G0435 Also See Notes	Counseling covered during preventive medicine services or a focused E/M visit. Labs as indicated.
Contraceptive methods and counseling (Recommendation 5.5)	The full range of Food and Drug Administration-approved contraceptive methods, sterilization procedures, and patient education and counseling for women with reproductive capacity.	11976, 11981, 11982, 11983, 57170, 58300, 58301, 58565, 58600, 58605, 58611, 58615, 58670, 58671, A4264, A4266, A4268, A4269, J1050, J7297, J7298, J7300, J7301, J7304, J7307, S4989	Counseling covered during preventive medicine services or a focused E/M visit. Oral contraceptives are included as a pharmacy benefit.

Michigan Department of Health and Human Services
Preventive Services Coverage Guidelines
Healthy Michigan Plan

Health and Medicine Division of the National Academies of Sciences, Engineering and Medicine

Recommended Preventive Health Care Services for Women

TYPE OF PREVENTIVE SERVICE	DESCRIPTION	RECOMMENDED ASSOCIATED CPT CODES-Medicaid Covered	NOTES
Breastfeeding support, supplies and counseling (Recommendation 5.6)	Comprehensive lactation support and counseling and costs of renting breastfeeding equipment. A trained provider should provide counseling services to all pregnant women and to those in the postpartum period to ensure the successful initiation and duration of breastfeeding. (The ACA ensures that breastfeeding counseling is covered; however, the committee recognizes that interpretation of this varies.)	99402-MIHP, A4281-A4286, E0602-E0604 S9443 Also See Notes	Lactation support and counseling is provided as part of childbirth education. It is covered during preventive medicine services or a focused E/M visit. IBCLC services may be provided as a benefit within the MIHP program or separately reimbursed in the outpatient setting when rendered by a qualified provider.
Screening and counseling for interpersonal and domestic violence (Recommendation 5.7)	Screening and counseling for interpersonal and domestic violence. Screening and counseling involve elicitation of information from women and adolescents about current and past violence and abuse in a culturally sensitive and supportive manner to address current health concerns about safety and other current or future health problems.	See Notes	Covered during preventive medicine services or a focused E/M visit.
Well-woman visit (Recommendation 5.8)	At least one well-women preventive care visit annually for adult women to obtain the recommended preventive services, including preconception and prenatal care. The committee also recognizes that several visits may be needed to obtain all necessary recommended preventive services, depending on a woman's health status, health needs, and other risk factors.	See Notes	Covered during preventive medicine services.

Early and Periodic Screening, Diagnosis and Treatment (EPSDT) services (19 and 20 year olds only)

Follow current Medicaid policy for coverage of EPSDT services.

Michigan Department of Health and Human Services

Preventive Services Coverage Guidelines

Healthy Michigan Plan

Revision History

04/2014	<ul style="list-style-type: none"> - Added HCPCS codes: A4281-A4286 and E0602-E0604, to IOM's Breastfeeding support, supplies and counseling preventive service. - Gestational diabetes mellitus screening added to USPSTF January 2014
07/2014	<ul style="list-style-type: none"> - Lung cancer screening added to USPSTF December 2013 - Hepatitis B screening: nonpregnant adolescents and adults added to USPSTF May 2014 - BRCA screening, counseling about had an update to the USPSTF preventive service title. - Dental caries prevention: preschool children had an update to the USPSTF preventive service title. - Align HIV screening with current 2014 USPSTF version.
10/2014	<ul style="list-style-type: none"> - Healthy diet counseling had an update to the USPSTF preventive service title and description. - Preeclampsia prevention: aspirin added to USPSTF September 2014 - Chlamydia screening: women had an update to the USPSTF preventive service title and description. - Gonorrhea screening: women had an update to the USPSTF description. - Sexually transmitted infections counseling had an update to the USPSTF description.
02/2015	<ul style="list-style-type: none"> - Added 2015 HCPCS codes: 45346, 45388, 77063, 77085-77086, 87623-87625, 87806, G0464, G0472, G6019, G6022, G6024 - Deleted discontinued HCPCS codes: 45339, 45383, 87620-87622
10/2015	<ul style="list-style-type: none"> - High blood pressure in adults: screening added to USPSTF October 2015 - Diabetes Screening had an update to the USPSTF description.
01/2016	<ul style="list-style-type: none"> - Added HCPCS code: A4264 - Added 2016 HCPCS codes: 80081, 81528, G0297, J7297, J7298 - Deleted HCPCS code: G0464 as it is a Grade I. - Deleted discontinued HCPCS codes: G6019, G6022, G6024, J7302
04/2016	<ul style="list-style-type: none"> - Depression screening: adults had an update to the USPSTF preventive service description. - Depression screening: adolescents had an update to the USPSTF preventive service description. - Added HCPCS code: S9443 - Revised notes concerning Lactation support.
07/2016	<ul style="list-style-type: none"> - Diabetes screening: Added HCPCS codes: 82952 - Hepatitis B screening: nonpregnant adolescents and adults: Deleted HCPCS codes: 86707, 87341 - Hepatitis B screening: pregnant women: Deleted HCPCS codes: 80055, 80081, 86707, 87341 - Osteoporosis screening: women: Deleted HCPCS codes: 77085 and 77086. Added HCPCS codes: 76977, 77078, 77081 - Syphilis screening: nonpregnant/pregnant persons had an update to the USPSTF preventive service description. - Aspirin to prevent cardiovascular disease: men/women had an update to the USPSTF preventive service title and description. - Blood pressure screening in adults had an update to the USPSTF preventive service description. - Tobacco use counseling: nonpregnant/pregnant had an update to the USPSTF preventive service description. - Anemia screening: pregnant women removed from USPSTF A and B recommendation. - Iron supplementation in children: removed from USPSTF A and B recommendation. - Colorectal cancer screening had an update to the USPSTF preventive service description.
11/2016	<ul style="list-style-type: none"> - Tuberculosis screening: adults added to USPSTF September 2016 - Breastfeeding counseling had an update to the USPSTF preventive service title and description. - Statin preventive medication: adults ages 40–75 years with no history of CVD, 1 or more CVD risk factors, and a calculated 10-year CVD event risk of 10% or greater added to USPSTF November 2016
01/2017	<ul style="list-style-type: none"> - Deleted discontinued HCPCS codes: G0436, G0437, 77052, 77057 - Added 2017 HCPCS codes: G0499, 77067 - Added HCPCS code: G0476